

AMENDMENT TO THE CLAIMS

Please cancel claims 43 to 85, without prejudice or disclaimer, and amend the claims as shown below in the listing of the claims.

1. (Original) A method of preparing nanoparticles having at least one polymer shell attached thereto comprising:
providing a type of nanoparticles; and
attaching a type of initiation monomers to the surfaces of the nanoparticles.
2. (Original) The method of Claim 1 wherein the initiation monomer comprises a cyclic olefin-containing group.
3. (Original) The method of Claim 2 wherein the initiation monomer comprises a norbornenyl group.
4. (Original) The method of Claim 1 wherein the nanoparticles are gold nanoparticles.
5. (Original) The method of Claim 4 wherein the initiation monomer is a norbornenyl- containing alkanethiol.
6. (Original) The method of Claim 5 wherein the initiation monomer is 1-mercapto-10-(*exo*-5-norbornen-2-oxy)-decane.
7. (Original) The method of Claim 1 wherein the initiation monomers are mixed with a type of attachment compounds, and both the initiation monomers and the attachment compounds are attached to the surfaces of the nanoparticles.
8. (Currently amended) The method of Claim 1 further comprising:
contacting the nanoparticles having the initiation monomers attached [to them] thereto with a transition metal ring-opening metathesis catalyst to activate the initiation monomers; and

contacting the nanoparticles with one or more types of propagation monomers of the formula P-L-N under conditions effective so that the propagation monomers are polymerized to form one or more polymer shells attached to the nanoparticles,

wherein:

N is a cyclic olefin-containing group;

P is a moiety which gives each polymer shell [a selected property or] one or more selected properties; and

L is a bond or linker whereby N is attached to P.

9. (Original) The method of Claim 8 wherein L is a polymer, ---COO--- , $\text{---CH}_2(\text{CH}_2)_m\text{COO---}$, ---OCO--- , $\text{---R}^1\text{N}(\text{CH}_2)_m\text{---NR}^1\text{---}$, $\text{---O}(\text{CH}_2)_m\text{---}$, $\text{---}(\text{CH}_2)_m\text{---}$, $\text{---N}(\text{R}^1)\text{C}(=\text{O})\text{---}(\text{CH}_2)_m\text{---}$, $\text{---O---}(\text{CH}_2)_m\text{---O---}$, $\text{---O---C}(=\text{O})\text{---}(\text{CH}_2)_m\text{---C}(=\text{O})\text{---O---}$, $\text{---C}(=\text{O})\text{---O---}(\text{CH}_2)_m\text{---O---C}(=\text{O})\text{---O---}$, $\text{---R}^1\text{N---}(\text{CH}_2)_m\text{---}$, $\text{---R}^1\text{N---C}(=\text{O})\text{---}(\text{CH}_2)_m\text{---C}(=\text{O})\text{---N---}$, $\text{---C}(=\text{O})\text{---N}(\text{R}^1)\text{---}(\text{CH}_2)_m\text{---N}(\text{R}^1)\text{---C}(=\text{O})\text{---}$, or $\text{---C}(=\text{O})\text{---N}(\text{R}^1)\text{---}(\text{CH}_2)_m\text{---}$, or comprises a moiety B that binds specifically to an analyte;

wherein:


R^1 has the formula $\text{X}(\text{CH}_2)_m$;

X is $-\text{CH}_3$, $-\text{CHCH}_3$, $-\text{COOH}$, $-\text{CO}_2(\text{CH}_2)_m\text{CH}_3$, $-\text{OH}$, $-\text{CH}_2\text{OH}$, ethylene glycol, hexa(ethylene glycol), $-\text{O}(\text{CH}_2)_m\text{CH}_3$, $-\text{NH}_2$, $-\text{NH}(\text{CH}_2)_m\text{NH}_2$, halogen, glucose, maltose, fullerene C_{60} , a cyclic olefin, or a nucleic acid; and

m is 0 – 30.

10. (Original) The method of Claim 8 wherein N is a norbornenyl-containing group.

11. (Original) The method of Claim 8 or 10 wherein the catalyst has the formula:



wherein:

M is osmium or ruthenium;

R^1 is hydrogen;

X^1 and X^2 , which may be different or the same, are any anionic ligand;

L^1 and L^2 , which may be different or the same, are any neutral electron donor;

and

R^2 is hydrogen, substituted or unsubstituted alkyl, or substituted or unsubstituted aryl.

12. (Original) The method of Claim 11 wherein M is ruthenium, R^1 is hydrogen, R^2 is phenyl, X^1 and X^2 are both $-\text{Cl}$, and L^1 and L^2 are both tricyclohexylphosphine.

13. (Currently amended) The method of Claim 8 or 10 wherein the catalyst has the formula:



wherein:

Re is rhenium (VII);

R¹ is selected from the group consisting of an alkyl having 1-20 carbon atoms, an aryl having 6-20 carbon atoms, an arralkyl having 7-30 carbon atoms, halogen substituted derivatives of [each] one of the alkyl, aryl, or arralkyl, and silicon-containing analogs of [each] one of the alkyl, aryl, or arralkyl;

R² is R¹ or is a substituent resulting from the reaction of the Re=CHR² moiety of the catalyst with an olefin that is being metathesized;

R³ and R⁴ are ligands which individually or together are sufficiently electron withdrawing to render the rhenium atom electrophilic enough for metathesis reaction; and

n is 1 or more.

14. (Original) The method of Claim 8 or 10 wherein the catalyst has the formula:



wherein:

M is molybdenum or tungsten;

R¹ and R² each individually may be an alkyl containing 1-20 carbon atoms, an aryl containing 6-20 carbon atoms, an arralkyl containing 7-20 carbon atoms, a halogen substituted derivative of the alkyl, aryl, or arralkyl, or a silicon-containing analog of one of the alkyl, aryl, or arralkyl; and

R³ is an alkyl containing 1-20 carbon atoms, an aryl containing 6-20 carbon atoms, an arralkyl containing 7-20 carbon atoms, or a substituent resulting from the reaction of the M=CHR³ moiety of said catalyst with an olefin being metathesized.

15. (Original) The method of Claim 8 or 10 wherein the nanoparticles are contacted with a single type of propagation monomers under conditions effective so that the monomers are polymerized to form a single polymer shell attached to the nanoparticles.

16. (Original) The method of Claim 15 wherein the polymer shell has redox activity.

17. (Original) The method of Claim 16 wherein the propagation monomer is *exo*-5-norbornen-2-yl ferrocenecarboxylate or *exo*-5-norbornen-2-yl ferroceneacetate.

18. (Original) The method of Claim 8 or 10 wherein:
the nanoparticles are contacted with a plurality of types of propagation monomers under conditions effective so that the monomers are polymerized to form one or more polymer shells attached to the nanoparticles, each polymer shell having one or more selected properties.
19. (Original) The method of Claim 18 wherein:
the nanoparticles are contacted with a first type of propagation monomers under conditions effective so that the monomers are polymerized to form a first polymer shell attached to the nanoparticles, the first polymer shell having a first selected property; and
then the nanoparticles are contacted with a second type of propagation monomers under conditions effective so that the monomers are polymerized to form a second polymer shell attached to the first polymer shell, the second polymer shell having a second selected property which is different from the first selected property of the first polymer shell.
20. (Original) The method of Claim 19 wherein one of the polymer shells has redox activity.
21. (Original) The method of Claim 20 wherein the propagation monomer polymerized to form the shell is *exo*-5-norbornen-2-yl ferrocenecarboxylate or *exo*-5-norbornen-2-yl ferroceneacetate.
22. (Original) The method of Claim 19 wherein the both polymer shells have redox activity.
23. (Original) The method of Claim 22 wherein the two polymer shells have different redox activities.
24. (Original) The method of Claim 23 wherein the propagation monomer polymerized to form the first polymer shell is *exo*-5-norbornen-2-yl ferrocenecarboxylate and the

propagation monomer polymerized to form the second polymer shell is *exo*-5-norbornen-2-yl ferroceneacetate.

25. (Original) The method of Claim 8 or 10 wherein the polymerization is stopped by adding a compound that terminates polymerization.

26. (Original) Nanoparticles having initiation monomers attached to them.

27. (Original) The nanoparticles of Claim 26 wherein the initiation monomers comprise cyclic olefin-containing groups.

28. (Original) The nanoparticles of Claim 27 wherein the initiation monomers comprise norbornenyl groups.

29. (Original) The nanoparticles of Claim 28 wherein the initiation monomers are norbornenyl-containing alkanethiols.

30. (Original) The nanoparticles of Claim 29 wherein the initiation monomers are 1-mercapto-10-(*exo*-5-norbornen-2-oxy)-decane.

31. (Original) Nanoparticles comprising one or more polymer shells attached to them, the polymer shells being formed by polymerizing one or more types of propagation monomers of the formula P-L-N,


wherein:


P is moiety which provides a desired property or properties to each of the polymer shells;

N is a cyclic olefin-containing group; and

L is a bond or a linker whereby N is attached to P.

32. (Currently amended) The nanoparticles of Claim 31 wherein L is a polymer,





wherein:

R^1 has the formula $\text{X}(\text{CH}_2)_m$;

X is $-\text{CH}_3$, $-\text{CHCH}_3$, $-\text{COOH}$, $-\text{CO}_2(\text{CH}_2)_m\text{CH}_3$, $-\text{OH}$, $-\text{CH}_2\text{OH}$, ethylene glycol, hexa(ethylene glycol), $-\text{O}(\text{CH}_2)_m\text{CH}_3$, $-\text{NH}_2$, $-\text{NH}(\text{CH}_2)_m\text{NH}_2$, halogen, glucose, maltose, fullerene C60, a cyclic olefin, or a nucleic acid; and

m is 0 – 30.

33. (Original) The nanoparticles of Claim 31 wherein N is a norbornenyl-containing group.
34. (Original) The nanoparticles of Claim 31 or 33 having a single polymer shell attached to them.

35. (Original) The nanoparticles of Claim 31 or 33 having a plurality of polymer shells attached to them.
36. (Original) The nanoparticles of Claim 35 having two polymer shells attached to them, the first polymer shell and the second polymer shell having different properties.
37. (Original) The nanoparticles of Claim 34 wherein the polymer shell has redox activity.
38. (Original) The nanoparticles of Claim 35 wherein one of the polymer shells has redox activity.
39. (Original) The nanoparticles of Claim 36 wherein the first polymer shell has redox activity and the second polymer shell has redox activity different than that of the first polymer shell.
40. (Original) The nanoparticles of Claim 31, 32, or 33 wherein a polymer shell comprises a binding moiety B that binds specifically to an analyte.
41. (Original) The nanoparticles of Claim 40 wherein the polymer shell comprising the binding moiety B is formed by polymerizing one or more types of binding monomers of the formula N-L-B, wherein N, L, and B have the same meanings as in Claim 40.
42. (Original) The nanoparticles of Claim 41 wherein the polymer shell comprising the binding moiety B is formed by polymerizing a mixture of one or more types of binding monomers and one or more types of propagation monomers.
43. to 85. (Currently cancelled)